



US006723549B2

(12) **United States Patent**
Miettinen-Oinonen et al.

(10) Patent No.: **US 6,723,549 B2**
(45) Date of Patent: ***Apr. 20, 2004**

(54) **CELLULASES, THE GENES ENCODING THEM AND USES THEREOF**

5,525,507 A 6/1996 Clarkson et al. 435/263
5,650,322 A 7/1997 Clarkson et al. 435/263
6,001,639 A 12/1999 Schulein et al. 435/263

(75) Inventors: **Arja Miettinen-Oinonen, Masala (FI); John Londesborough, Helsinki (FI); Jari Vehmaanperä, Klaukkala (FI); Heli Haakana, Rajamäki (FI); Arja Mäntylä, Helsinki (FI); Raija Lanto, Klaukkala (FI); Minna Elovainio, Helsinki (FI); Vesa Joutsjoki, Helsinki (FI); Marja Paloheimo, Vantaa (FI); Pirkko Suominen, Helsinki (FI)**

(73) Assignee: **AB Enzymes Oy (FI)**

(*) Notice: This patent issued on a continued prosecution application filed under 37 CFR 1.53(d), and is subject to the twenty year patent term provisions of 35 U.S.C. 154(a)(2).

Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **08/841,636**

(22) Filed: **Apr. 30, 1997**

(65) **Prior Publication Data**

US 2002/0168751 A1 Nov. 14, 2002

Related U.S. Application Data

(63) Continuation of application No. PCT/FI96/00550, filed on Oct. 17, 1996, and a continuation-in-part of application No. 08/732,181, filed on Oct. 16, 1996.

(60) Provisional application No. 60/005,335, filed on Oct. 17, 1995, provisional application No. 60/007,926, filed on Dec. 4, 1995, and provisional application No. 60/020,840, filed on Jun. 28, 1996.

(51) Int. Cl.⁷ **C12N 9/42; C11D 3/386**

(52) U.S. Cl. **435/204; 435/263; 510/392**

(58) Field of Search **475/209, 263; 510/392**

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,438,864 A 4/1969 Tendler 195/66
3,664,961 A 5/1972 Norris 252/99
4,106,989 A 8/1978 Komura et al. 195/62
4,832,864 A 5/1989 Olson 252/174.12
4,894,338 A 1/1990 Knowles et al. 435/172.3
4,912,056 A 3/1990 Olson 435/263
5,006,126 A 4/1991 Olson et al. 8/401
5,120,463 A 6/1992 Bjork et al. 252/174.12
5,122,159 A 6/1992 Olson et al. 8/401
5,213,581 A 5/1993 Olson et al. 8/401
5,232,851 A 8/1993 Cox et al. 435/263
5,246,853 A 9/1993 Clarkson et al. 435/263
5,290,474 A 3/1994 Clarkson et al. 252/174.12
5,298,405 A 3/1994 Nevalainen et al. 435/209
5,443,750 A 8/1995 Convents et al. 252/174.12
5,520,838 A 5/1996 Baeck et al. 252/174.12

FOREIGN PATENT DOCUMENTS

DE 1 767 737 9/1971
EP 0 220 016 4/1987
EP 0 238 216 9/1987
EP 0 307 564 3/1989
EP 0 383 828 8/1990
EP 0 406 314 1/1991
EP 0 510 091 10/1992
EP 0 521 999 1/1993
EP 0 531 372 3/1993
EP 0 576 526 1/1994
GB 2 075 028 11/1981
WO WO 89/09259 10/1989
WO WO 91/14822 10/1991
WO WO 91/17243 11/1991
WO WO 91/17244 11/1991
WO WO 92/06165 4/1992
WO WO 92/06209 4/1992
WO WO 92/06210 4/1992
WO WO 92/06221 4/1992
WO WO 92/17572 10/1992
WO WO 92/17574 10/1992
WO WO 92/18599 10/1992
WO WO 93/11249 6/1993
WO WO 93/22428 11/1993
WO WO 93/24621 12/1993
WO WO 93/24622 12/1993
WO WO 94/07983 4/1994
WO WO 94/07998 4/1994
WO WO 94/14953 7/1994
WO WO 94/23113 10/1994
WO WO 95/02675 1/1995
WO WO 95/16782 6/1995
WO WO 95/24471 9/1995
WO WO 95/25840 9/1995
WO WO 96/17994 6/1996
WO WO 96/29397 9/1996
WO WO 96/34945 11/1996

OTHER PUBLICATIONS

Sequence search, 1999.*

(List continued on next page.)

Primary Examiner—Charles L. Patterson, Jr.

(74) Attorney, Agent, or Firm—Sterne, Kessler, Goldstein & Fox P.L.L.C.

(57) **ABSTRACT**

Genes encoding novel cellulases, and a gene encoding a protein that facilitates the action of such novel cellulases, the novel cellulases and a protein that facilitates the action of such cellulases, and enzyme preparations containing such proteins are described. The native hosts and the culture medium of said hosts containing said novel cellulases are also disclosed. These proteins are especially useful in the textile and detergent industry and in pulp and paper industry.

16 Claims, 37 Drawing Sheets

U.S. patent
Seq'd NO: 31.
NO ODP. or DP.

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

CCGCGGACTG GCATC

15

(2) INFORMATION FOR SEQ ID NO:45:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 16 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

CCGCGGACTG CGCATC

16

What is claimed is:

1. An isolated and essentially homogenous polypeptide as analyzed on SDS-PAGE having the amino acid sequence set forth in FIG. 19 (SEQ ID NO:31).

2. An isolated and essentially homogenous polypeptide as analyzed on SDS-PAGE having an amino acid sequence which has at least 80% identity to the amino acid sequence set forth in FIG. 19 (SEQ ID NO:31).

3. An enzyme preparation having cellulase activity, wherein said activity comprises the cellulase activity of a 20K polypeptide having the amino acid sequence set forth in FIG. 19 (SEQ ID NO:31), wherein said enzyme preparation is obtained by a process comprising:

- (i) culturing a host cell transformed with the nucleic acid sequence set forth in FIG. 19 (SEQ ID NO:30); and
- (ii) separating said host cell from the culture medium and obtaining the supernatant having said 20K cellulase activity;

wherein the level of 20K cellulase activity in the supernatant obtained from said transformed host cell is at least three times the level of 20K cellulase activity obtained from the supernatant of the untransformed host cell naturally expressing said cellulase when said untransformed host cell is cultivated under the same conditions.

4. An enzyme preparation having cellulase activity wherein said enzyme preparation is obtained by a process comprising:

- (i) culturing a host cell transformed with a nucleic acid sequence encoding a cellulase having at least 80% identity to the amino acid sequence of the 20K cellulase set forth in FIG. 19 (SEQ ID NO:31); and
- (ii) separating said host cell from the culture medium and obtaining the supernatant having said 20K cellulase activity or cellulase activity of a cellulase that has at least 80% identity to the sequence of said 20K cellulase set forth in FIG. 19 (SEQ ID NO:31),

wherein the level of said 20K cellulase activity or the level of said activity of said cellulase that has at least 80% identity to said sequence of said 20K cellulase set forth in FIG. 19 (SEQ ID NO: 31) in the supernatant obtained from said transformed host cell is at least three times the level of said 20K cellulase activity or activity of said cellulase that has at least 80% identity to said sequence of said 20K cellulase set forth in FIG. 19 (SEQ ID NO:31), respectively, obtained from the supernatant of the strain naturally expressing said cellulase when said strain is cultivated under the same conditions.

5. An enzyme preparation comprising at least one cellulase from *Melanocarpus albomyces* CBS 685.95.

6. The enzyme preparation of any one of claims 3-5, which is in the form of a liquid.

7. The enzyme preparation of any one of claims 3-5, which is in the form of a dry powder.

8. A detergent composition comprising the enzyme preparation of any one of claims 3-5 in combination with a surface active agent or surfactant.

9. A method of treating cellulosic fiber containing textile material, said method comprising contacting said textile material with the detergent composition of claim 8.

10. An isolated and essentially homogenous polypeptide as analyzed on SDS-PAGE having amino acids 22-235 of the amino acid sequence set forth in FIG. 19 (SEQ ID NO:31).

11. An isolated and essentially homogenous polypeptide as analyzed on SDS-PAGE having an amino acid sequence which has at least 80% identity to amino acids 22-235 of the amino acid sequence set forth in FIG. 19 (SEQ ID NO:31).

12. An enzyme preparation having cellulase activity, wherein said activity comprises the cellulase activity of a polypeptide having amino acids 22-235 of the amino acid sequence set forth in FIG. 19 (SEQ ID NO:31), wherein said enzyme preparation is obtained by a process comprising:

- (i) culturing a host cell transformed with a nucleic acid sequence encoding amino acids 22-235 of the amino acid sequence set forth in FIG. 19 (SEQ ID NO:31); and

(ii) separating said host cell from the culture medium and obtaining the supernatant having said cellulase activity; wherein the level of said cellulase activity in the supernatant obtained from said transformed host cell is at least three times the level of said cellulase activity obtained from the supernatant of the strain naturally expressing said cellulase when said strain is cultivated under the same conditions.

13. An enzyme preparation having cellulase activity wherein said enzyme preparation is obtained by a process comprising:

- (i) culturing a host cell transformed with a nucleic acid sequence encoding a cellulase having at least 80% identity to amino acids 22-235 of the amino acid sequence set forth in FIG. 19 (SEQ ID NO:31); and

(ii) separating said host cell from the culture medium and obtaining the supernatant having the cellulase activity of a cellulase that has at least 80% identity to amino acids 22-235 of the amino acid sequence set forth in FIG. 19 (SEQ ID NO:31);

95

wherein the level of said activity of said cellulase that has at least 80% identity to amino acids 22-235 of the amino acid sequence set forth in FIG. 19 (SEQ ID NO:31) in the supernatant obtained from said transformed host cell is at least three times the level of said activity of said cellulase that has 80% identity to amino acids 22-235 of the amino acid sequence set forth in FIG. 19 (SEQ ID NO:31) obtained from the supernatant of the strain naturally expressing said cellulase when said strain is cultivated under the same conditions.

96

14. The enzyme preparation of either claim 12, or claim 13, which is in the form of a liquid.

15. The enzyme preparation of either claim 12 or claim 13, which is in the form of a dry powder.

16. A detergent composition comprising the enzyme preparation of either claim 12 or claim 13 in combination with a surface active agent or surfactant.

* * * * *